

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	11.5 (9.2-14.4)	1,411	14.1 (9.7-20.1)	248			●
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	22.0 (18.1-26.5)	1,404	30.2 (22.4-39.3)	246		●	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	4.4 (2.2-8.7)	632	10.9 (5.6-20.1)	89			●
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	26.5 (21.9-31.8)	670	31.3 (19.7-45.9)	98			●
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	10.0 (7.9-12.5)	1,390	17.4 (11.6-25.2)	237		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	2.2 (1.4-3.5)	1,386	6.4 (3.2-12.4)	234		●	
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)	5.6 (3.8-8.2)	1,409	11.6 (7.3-17.8)	250		●	
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	7.8 (6.1-10.0)	1,411	14.6 (8.4-24.1)	248			●
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	5.9 (4.2-8.2)	1,407	10.8 (5.6-20.1)	248			●
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	22.3 (19.0-25.9)	1,378	32.8 (26.8-39.5)	233		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	8.6 (6.3-11.6)	1,373	16.6 (11.7-22.9)	232		●	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	6.5 (5.1-8.2)	1,389	13.9 (9.9-19.2)	233		●	
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)	12.2 (10.2-14.5)	1,359	22.7 (15.9-31.3)	235		●	
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	3.1 (2.1-4.4)	867	8.8 (4.6-16.2)	145			●
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	6.3 (5.0-8.0)	901	14.7 (9.4-22.2)	160		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	13.6 (11.1-16.6)	1,382	22.4 (16.4-29.7)	239		●	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	9.7 (7.6-12.3)	1,405	21.4 (15.1-29.3)	241		●	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	29.9 (27.4-32.4)	1,388	59.9 (52.1-67.2)	238		●	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	13.7 (11.7-16.0)	1,395	38.1 (31.6-45.0)	238		●	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	10.9 (9.2-13.0)	1,395	35.6 (30.0-41.6)	233		●	
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	8.3 (6.1-11.2)	1,144	28.4 (22.3-35.5)	197		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
Q29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	3.8 (2.3-6.0)	1,147	10.7 (6.2-18.1)	195		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)	25.3 (21.8-29.1)	1,293	34.1 (27.7-41.3)	206		●	
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	7.8 (6.3-9.5)	1,300	14.5 (9.0-22.4)	217			●
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	4.5 (3.1-6.6)	1,335	11.7 (7.0-18.8)	220		●	
QNFRCIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	0.8 (0.4-1.7)	1,335	1.8 (0.5-6.2)	220			●
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.5 (0.2-1.4)	1,335	1.6 (0.4-6.1)	220			●
QN33: Percentage of students who smoked more than 10 cigarettes per day (on the days they smoked during the 30 days before the survey, among students who currently smoked cigarettes)	6.1 (1.5-21.9)	59	-	28			

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in the denominator (subgroup).

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])	35.3 (32.0-38.8)	1,349	42.6 (36.9-48.5)	223		●	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	5.1 (3.2-8.0)	1,241	12.5 (7.6-20.0)	201		●	
QN36: Percentage of students who usually got their own electronic vapor products by buying them in a store (such as a convenience store, supermarket, discount store, gas station, or vape store, during the 30 days before the survey, among students who currently used electronic vapor products and who were aged <18 years)	14.9 (7.9-26.4)	46	-	23			
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, Copenhagen, Camel Snus, Marlboro Snus, General Snus, Ariva, Stonewall, or Camel Orbs], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	2.3 (1.6-3.4)	1,376	10.6 (5.1-20.8)	245		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in the denominator (subgroup).

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	5.2 (3.6-7.5)	1,399	12.5 (6.8-21.8)	244		●	
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.8 (5.7-10.5)	1,339	18.8 (11.4-29.3)	231		●	
QNTB3: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	8.8 (6.5-11.8)	1,324	20.3 (12.0-32.2)	232		●	
QNTB4: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	11.2 (8.4-14.9)	1,220	24.7 (15.4-37.0)	210		●	
QN39: Percentage of students who did not try to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)	57.3 (49.3-64.9)	233	40.7 (28.4-54.2)	59	●		

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN40: Percentage of students who ever drank alcohol (at least one drink of alcohol, on at least 1 day during their life)	54.4 (49.1-59.7)	1,264	73.7 (62.6-82.4)	211		●	
QN41: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	15.0 (12.7-17.7)	1,257	25.5 (17.2-36.0)	213		●	
QN42: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	21.3 (17.4-25.9)	1,195	36.8 (26.7-48.3)	191		●	
QN43: Percentage of students who usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	42.9 (37.4-48.7)	255	49.9 (40.1-59.7)	68			●
QN44: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	8.5 (6.3-11.4)	1,302	12.5 (8.2-18.6)	207			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN45: Percentage of students who reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours, during the 30 days before the survey)	1.6 (0.9-2.7)	1,154	0.0 (.-)	180	●		
QN46: Percentage of students who ever used marijuana (one or more times during their life)	41.9 (37.5-46.5)	1,345	58.1 (50.8-65.0)	217		●	
QN47: Percentage of students who tried marijuana for the first time before age 13 years	6.6 (5.1-8.4)	1,360	15.2 (9.6-23.3)	224		●	
QN48: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	22.4 (19.3-25.9)	1,375	36.8 (30.1-44.0)	229		●	
QN49: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	4.4 (3.1-6.2)	1,394	13.4 (6.6-25.3)	249		●	
QN50: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	7.8 (5.9-10.4)	1,408	16.4 (10.4-24.8)	249		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN51: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	2.5 (1.4-4.5)	1,406	11.7 (6.3-20.8)	248		●	
QN52: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	2.4 (1.5-4.0)	1,392	10.5 (4.7-21.9)	246		●	
QN53: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	3.7 (2.2-6.4)	1,411	12.5 (6.6-22.5)	249		●	
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)	2.9 (1.8-4.7)	1,410	14.7 (7.9-25.8)	248		●	
QN56: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	12.9 (10.6-15.5)	1,371	21.9 (15.4-30.2)	244		●	
QN57: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	2.1 (1.3-3.5)	1,404	12.2 (6.4-22.0)	245		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN58: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	31.6 (28.2-35.3)	1,394	33.7 (27.3-40.9)	237			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	38.8 (33.9-43.9)	1,186	46.5 (36.8-56.4)	167			●
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	5.1 (4.1-6.2)	1,189	7.1 (3.4-14.2)	172			●
QN61: Percentage of students who had sexual intercourse with four or more persons during their life	10.1 (7.6-13.4)	1,187	10.9 (6.1-18.7)	171			●
QN62: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	28.8 (24.1-33.9)	1,182	33.6 (23.5-45.5)	170			●
QN63: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	13.3 (8.9-19.4)	346	24.5 (15.2-37.1)	56		●	
QN64: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	44.6 (38.2-51.2)	331	55.7 (37.8-72.3)	52			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN65: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	86.1 (80.7-90.2)	336	81.6 (53.7-94.4)	53			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	96.6 (94.1-98.1)	336	98.9 (91.4-99.9)	53			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	93.0 (86.6-96.4)	336	96.3 (88.8-98.8)	53			●
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	75.7 (67.4-82.4)	336	76.7 (52.4-90.8)	53			●
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	94.0 (90.9-96.1)	331	95.3 (82.8-98.9)	49			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	17.0 (12.4-22.9)	336	27.4 (12.7-49.5)	53			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	17.4 (14.9-20.3)	1,281	18.1 (12.7-25.2)	228			●
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	17.4 (14.6-20.7)	1,281	21.0 (16.5-26.4)	228			●
QN68: Percentage of students who described themselves as slightly or very overweight	32.7 (29.3-36.3)	1,397	42.2 (33.2-51.8)	248			●
QN69: Percentage of students who were not trying to lose weight	50.9 (47.0-54.7)	1,392	45.4 (38.3-52.6)	247			●
QN70: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	25.4 (22.7-28.3)	1,380	31.2 (24.8-38.5)	247			●
QN71: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	10.5 (8.2-13.3)	1,386	16.2 (12.4-20.8)	247		●	

*Confidence interval.

[†]Based on t-test analyses, $p < 0.05$.

[§]Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	4.6 (3.3-6.4)	1,374	7.1 (4.1-12.1)	245			●
QN72: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	48.3 (44.8-51.8)	1,377	44.2 (37.6-51.0)	247			●
QN73: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)	43.1 (39.5-46.7)	1,373	38.4 (32.8-44.3)	247			●
QN74: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)	58.8 (55.2-62.4)	1,379	57.6 (47.3-67.4)	247			●
QN75: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)	27.5 (24.0-31.4)	1,378	29.5 (23.0-37.0)	247			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	12.5 (10.2-15.2)	1,357	10.3 (6.8-15.1)	245			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	17.1 (14.2-20.5)	1,355	20.5 (15.9-26.0)	242			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	11.6 (9.2-14.6)	1,355	14.6 (10.1-20.5)	242			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	6.0 (4.3-8.5)	1,355	8.7 (5.5-13.5)	242			●
QN77: Percentage of students who did not drink milk (during the 7 days before the survey)	24.2 (20.6-28.2)	1,361	35.0 (28.6-41.8)	243		●	
QN78: Percentage of students who did not eat breakfast (during the 7 days before the survey)	20.0 (17.4-22.8)	1,356	18.8 (13.9-24.9)	244			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	74.9 (71.9-77.6)	1,356	83.9 (76.5-89.3)	244		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	60.7 (56.4-64.8)	1,351	72.2 (63.5-79.6)	239		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	17.5 (14.8-20.4)	1,351	23.7 (18.4-30.0)	239		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	81.4 (78.6-84.0)	1,351	87.8 (80.6-92.6)	239			●
QN80: Percentage of students who watched television 3 or more hours per day (on an average school day)	23.4 (20.0-27.3)	1,358	23.3 (17.5-30.4)	241			●
QN81: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	40.8 (38.2-43.6)	1,350	39.2 (30.8-48.3)	242			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN82: Percentage of students who did not attend physical education (PE) classes on 1 or more days (in an average week when they were in school)	30.2 (23.5-38.0)	1,327	34.1 (25.7-43.6)	240			●
QNDLYPE: Percentage of students who did not attend physical education classes on all 5 days (in an average week when they were in school)	56.1 (45.3-66.4)	1,327	66.2 (55.2-75.8)	240		●	
QN83: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	48.4 (44.7-52.2)	1,341	52.5 (45.0-59.8)	241			●
QN88: Percentage of students who did not get 8 or more hours of sleep (on an average school night)	82.9 (80.9-84.7)	1,318	84.6 (76.9-90.0)	241			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	81.2 (78.3-83.7)	1,299	77.3 (71.1-82.5)	232			●
QN86: Percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)	29.9 (26.7-33.3)	1,346	40.0 (33.8-46.6)	242		●	
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)	1.8 (1.2-2.8)	1,346	2.3 (0.7-7.4)	242			●
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	21.0 (18.4-23.8)	1,309	31.3 (25.5-37.8)	233		●	
QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	29.6 (25.3-34.4)	1,306	28.0 (21.0-36.3)	239			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN90: Percentage of students who have ever been hit, beaten, kicked, or physically hurt in any way by an adult (not including being spanked for bad behavior)	16.0 (13.2-19.3)	1,385	24.8 (19.4-31.1)	237		●	
QN91: Percentage of students who have ever seen or heard adults in their home slap, hit, kick, punch, or beat each other up	17.2 (14.6-20.3)	1,401	29.4 (21.2-39.2)	238		●	
QN92: Percentage of students who have been the victim of teasing or name calling because someone thought they were gay, lesbian, or bisexual (during the 12 months before the survey)	5.5 (4.3-7.0)	1,396	23.4 (16.7-31.7)	239		●	
QN93: Percentage of students who have been pregnant or gotten someone pregnant (one or more times)	3.5 (2.5-4.8)	1,401	10.6 (4.6-22.4)	245			●
QN94: Percentage of students who did not see a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	27.9 (24.2-31.9)	1,324	32.5 (24.2-42.1)	240			●
QN95: Percentage of students who preported their doctor or nurse had not discussed ways to prevent pregnancy, AIDS, or other sexually transmitted diseases (STDs) or they were not sure if they had (during their last check-up, among students who have had a check-up)	51.5 (47.9-55.2)	1,134	51.4 (44.4-58.3)	190			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Chicago High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who have never had or were not sure if they have ever had the HPV vaccine, a vaccine to prevent human papillomavirus or HPV infection (also called the cervical cancer vaccine, HPV shot, or GARDASIL)	64.6 (60.9-68.1)	1,311	48.5 (41.7-55.4)	233	●		
QN97: Percentage of students who usually slept in the home of a friend, family member, or other person because they had to leave their home or their parent or guardian cannot afford housing (during the 30 days before the survey)	4.1 (3.0-5.8)	1,312	9.5 (5.2-16.6)	241		●	
QN98: Percentage of students who have ever slept away from their parents or guardians because they were kicked out, ran away, or were abandoned (during the 30 days before the survey)	9.9 (7.7-12.5)	1,286	16.9 (8.7-30.4)	228			●
QN100: Percentage of students who have not had or are not sure if they have had sex education in school	16.4 (12.3-21.4)	1,293	22.4 (17.1-28.8)	235			●
QN101: Percentage of students who reported there is not at least one teacher or other adult in their school that they can talk to if they have a problem or who were not sure if there is at least one teacher or other adult in their school that they can talk to if they have a problem	24.7 (21.3-28.5)	1,300	29.1 (21.5-38.1)	238			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.